



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ing, rearing, and treatment of Pheasants in captivity. A dozen species and several interesting hybrids are beautifully illustrated in the 16 fine plates that accompany the text, one of which shows the Argus Pheasant displaying its plumage. The principal subjects treated are the 'Natural History of Pheasants,' 'Management in Preserves,' 'Management in Confinement,' 'Diseases of Pheasants,' 'Pheasants adapted to the Covert,' and 'Pheasants adapted to the Aviary.' No species of Pheasant seems susceptible to domestication, like the common fowl; even when bred in aviaries for generations they retain their original wildness, and when set at liberty "betake themselves to the woods and coverts as soon as able to shift for themselves." Mr. Tegetmeier refers to the successful introduction of the Ring-necked and Golden Pheasants into Oregon, where they have become thoroughly acclimatized; but he seems to take a too sanguine view of the case of the English Pheasant, introduced in the Eastern States, or to have been misinformed, when he states it "is generally regarded as the future game bird of the country," etc. (p. 38)—J. A. A.

Bird-nesting with a Camera.¹—The second part² of Mr. Lee's admirable work is fully up to the standard of excellence set by Part I,—higher praise cannot be accorded it.

The nests of the following species are herein figured and described:

(1) Coot (*Fulica atra*), two plates; (2) Shoveler (*Spatula clypeata*); (3) Kittiwake Gull (*Rissa tridactyla*); (4) Waterhen (*Gallinula chloropus*), two plates; (5) Willow Wren (*Phylloscopus trochilus*); (6) Sandwich Tern (*Sterna cantiaca*), and (7) Sheld-Duck (*Tadorna cornuta*), two plates.—F. M. C.

A List of the Birds of Florida.³—This is a nominal list of the birds of Florida, enumerating 352 species and subspecies, and will doubtless prove a convenient Check-List to students of Florida birds. The following species have apparently been overlooked: *Buteo borealis harlani* (see Bendire, Life Histories of N. A. Birds, Part I, p. 218); *Spiza americana* (see Beckham, Bull. N. O. C., VII, 1882, p. 250); *Dendroica dominica albilora* (see Scott, Auk, XII, 1890, p. 20); *Sylvania canadensis* (see Allen, Bull. Mus. Comp. Zoöl. II, 1891, p. 269); *Seiurus noveboracensis notabilis* (see Brewster and Chapman, Auk, VIII, 1891, p. 135).—F. M. C.

¹ Among British Birds in their Nesting Haunts. Illustrated by the Camera. By Oswin A. J. Lee. Part II. Edinburgh. David Douglas. Folio, pp. 41-77, pll. x.

² For a notice of Part I, see this journal, Vol. XIV, 1897, p. 106.

³ A List | of the | Birds of Florida | By Charles B. Cory | For sale by | Bradly Whidden | 18 Arch St. Boston. | 1896—8vo, pp. 24.